Carbohydrate Research, 245 (1992) C9–C11
Elsevier Science Publishers B.V., Amsterdam

CONTENTS

Avoiding oxidative degradation during sodium hydroxide/methyl iodide-mediated carbohydrate methylation in dimethyl sulfoxide
P.W. Needs and R.R. Selvendran (Norwich, United Kingdom) 1

Proton and carbon NMR chemical-shift assignments for [β-D-Fru*f*-(2 → 1)]$_3$-(2 ↔ 1)-α-D-Glc*p* (nystose) and [β-D- Fru*f*-(2 → 1)]$_4$-(2 ↔ 1)-α-D-Glc*p* (1,1,1-kestopentaose) from two-dimensional NMR spectral measurements
J. Liu, A.L. Waterhouse (Davis, CA, USA) and N.J. Chatterton (Logan, UT, USA) 11

Conformational analysis of methyl β-cellobioside by ROESY NMR spectroscopy and MD simulations in combination with the CROSREL method
L.M.J. Kroon-Batenburg, J. Kroon, B.R. Leeflang and J.F.G. Vliegenthart (Utrecht, Netherlands) 21

Interaction of 8-anilino-1-naphthalenesulfonate with 2,6-di-*O*-methylcyclomaltoheptaose
J. Nishijo, M. Nagai and M. Yasuda (Kobe, Japan) 43

The search for D-glucose derivatives suitable for the study of natural hydrogen isotope fractionation
J.M. Williams, H. Jegou, V. Langlois (Nantes, France), B.-L. Zhang and M.L. Martin (Swansea, United Kingdom) 57

Synthesis of *p*-trifluoroacetamidophenyl 6-deoxy-2-*O*-{3-*O*-[2-*O*-methyl-3-*O*-(2-*O*-methyl-α-D-rhamnopyranosyl)- α-L-fucopyranosyl]-α-L-rhamnopyranosyl}-α-L-talopyranoside: a spacer armed tetrasaccharide glycopeptidolipid antigen of *Mycobacterium avium* serovar 20
J. Kerékgyártó, Z. Szurmai and A. Lipták (Debrecen, Hungary) 65

Synthesis of the laminara-oligosaccharide methyl β-glycosides of dp 3–8
K. Takeo, K. Maki, Y. Wada and S. Kitamura (Kyoto, Japan) 81

Interpretation of dextransucrase inhibition at high sucrose concentrations
A. Tanriseven and J.F. Robyt (Ames, IA, USA) 97

Structural studies of the antigenic polysaccharide of *Eubacterium saburreum*, strain T19
N. Sato, F. Nakazawa, M. Sato, E. Hoshino (Niigata, Japan) and T. Ito (Tokyo, Japan) .. 105

NMR studies of oligosaccharides derived from hyaluronate: complete assignment of ^{1}H and ^{13}C NMR spectra of aqueous di- and tetra-saccharides, and comparison of chemical shifts for oligosaccharides of increasing degree of polymerisation
R. Toffanin, B.J. Kvam, A. Flaibani, M. Atzori, F. Biviano (Abano Terme, Italy) and S. Paoletti (Trieste, Italy) 113

A polysaccharide produced by a mucoid strain of *Moraxella nonliquefaciens* with a 2-acetamido-2-deoxy-5-*O*-(3-deoxy-β-D-*manno*-octulopyranosyl)-β-D-galactopyranosyl repeating unit
R. Reistad (Lysaker, Norway), U. Zähringer (Borstel, Germany), K. Bryn, J. Alstad, K. Bøvre and E. Jantzen (Oslo, Norway) 129

Chemical synthesis and kinetic characterization of UDP-2-deoxy-D-lyxo-hexose("UDP-2-deoxy-D-galactose"), a donor-substrate for β-(1 → 4)-D-galactosyltransferase
G. Srivastava, O. Hindsgaul and M.M. Palcic (Alberta, Canada) 137

Notes

Synthesis and characterization of two 1,2,3,6,2′,3′,4′,6′-octa-*O*-benzoyl-β-D-hexapyranosyl-(1 → 4)β-D-allopyranoses
N.B. D'Accorso and I.M.E. Thiel (Buenos Aires, Argentina) 145

Isolation and purification of trehalose 6-mono- and 6,6′-di-corynomycolates from *Corynebacterium matruchotii*. Structural characterization by ^{1}H NMR
A.K. Datta and K. Takayama (Madison, WI, USA) 151

Isolation and structural analysis of the tetrasaccharide 3-deoxy-5-*O*-[3-*O*-(3-*O*-α-D-glucopyranosyl-L-*glycero*-α-D-*manno*-heptopyranosyl)-L-*glycero*-α-D-*manno*-heptopyranosyl]-D-*manno*-octulosonic acid from the core region of the lipopolysaccharide of *Salmonella minnesota* strain R5 (rough mutant chemotype RcP$^-$)
O. Holst and H. Brade (Borstel, Germany) 159

Regular papers

An equilibrium study of *p*-Methyl Red inclusion complexes with α- and β-cyclodextrins
K.M. Tawarah and S.J. Khouri (Irbid, Jordan) 165

Synthesis of multivalent β-lactosyl clusters as potential tumor metastasis inhibitors
B. Dean, H. Oguchi, S. Cai, E. Otsuji, K. Tashiro, S.-i. Hakomori and T. Toyokuni (Washington, DC, USA) 175

Kohlenhydrat-Bausteine zum Aufbau Tumor-assoziierter Antigene. Synthese von Lewis Y-Determinanten
W. Kinzy und A. Löw (Darmstadt, Deutschland) 193

Di- and tri-saccharide glycosyl donors for the synthesis of fragments of the O-specific antigen of *Shigella dysenteriae* type 1
P. Kováč (Bethesda, Maryland, USA) 219

Preparation of 4,6-cyclo-4,6-dideoxy-hexopyranoses by palladium-mediated intramolecular cyclodehalogenation
H.P. Wessel, M.-C. Viaud and V. Gardon (Basel, Switzerland) 233

Convergent synthesis of an elusive hexasaccharide corresponding to the cell-wall polysaccharide of the β-hemolytic *Streptococcus* Group A
J.-R. Marino-Albernas, S.L. Harris, V. Varma and B.M. Pinto (Burnaby, Canada) 245

On the structure of calonyctin A, a plant growth regulator
Y. Fang, W. Chai, S. Chen, Y. He, L. Zhao, J. Peng, H. Huang and B. Xin (Beijing, China) 259

Structure of an extracellular polysaccharide produced by *Erwinia chrysanthemi*
J.S.S. Gray, J.M. Brand (Alice, South Africa), T.A.W. Koerner (Iowa City, IA, USA) and R. Montgomery (Alice, South Africa) 271

Resistance to acid hydrolysis of lipid-complexed amylose and lipid-free amylose in lintnerised waxy and non-waxy barley starches
W.R. Morrison, R.F. Tester (Glasgow, United Kingdom), M.J. Gidley (Bedford, United Kingdom) and J. Karkalas (Glasgow, United Kingdom) 289

Structural studies of acetan, an exopolysaccharide elaborated by *Acetobacter xylinum*
P.-E. Jansson, J. Lindberg, K.M.S. Wimalasiri (Stockholm, Sweden) and M.A. Dankert (Buenos Aires, Argentina) 303

Structural studies of the capsular polysaccharide (S-21) from *Klebsiella pneumoniae* ATCC 31314
P. Edebrink, P.-E. Jansson and G. Widmalm (Stockholm, Sweden) 311

Recognition of the acceptor β-D-Glc*p*NAc-(1 → 2)-α-D-Man*p*-(1 → 6)-β-D-Glc*p*-OR by *N*-acetylglucosaminyltransferase-V: None of the hydroxyl groups on the Glc-residue are important
T. Linker, S.C. Crawley and O. Hindsgaul (Alberta, Canada) 323

Notes

Complete assignment of the ^{1}H NMR spectra of phytoalexin elicitor-active oligoglucosides
V.-M. Ló, M.G. Hahn (Athens, Georgia, USA), N. Hong, T. Ogawa (Saitama, Japan) and H. Van Halbeek (Athens, Georgia, USA) 333

Synthetic mirror cord factors: Protected dimycolyl esters of an α,α-(1 ↔ 1)-bisheptosiduronic acid
H.H. Baer and X. Wu (Ottawa, Canada) 347

Identification of oligosaccharides consisting of D-glucuronic acid and L-*glycero*-D-*manno*- and D-*glycero*-D-*manno*-heptose isolated from *Vibrio parahaemolyticus* O2 lipopolysaccharide
S. Kondo, T. Watabe (Saitama, Japan), Y. Haishima (Tokyo, Japan) and K. Hisatsune (Saitama, Japan) 353

Occurrence of 2-*O*-sulphated D-glucuronic acid in rat liver heparan sulphate
J. Kovensky and A.F. Cirelli (Buenos Aires, Argentina) 361